

SP-D

Catalog # PVGS1546

Product Information

Primary Accession P35247
Species Human

Sequence Ala21-Phe375

Purity > 98% as analyzed by SDS-PAGE

Endotoxin Level

Expression System HEK 293

Formulation Lyophilized from a 0.2 Im filtered solution in PBS.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH₂O or PBS up to 100 □g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Gene ID 6441

Other Names Pulmonary surfactant-associated protein D, PSP-D, SP-D, Collectin-7, Lung

surfactant protein D, SFTPD, COLEC7, PSPD, SFTP4

 Target Background
 Surfactant protein D is a member of the collectin family of innate immune

modulators. It is constitutively secreted by alveolar lining cells and epithelium associated with tubular structures, that contains a N-terminal collagen-like domain and a C-terminal lectin domain that are characteristic of members of the collectin family of proteins, and also shows calcium-dependent binding to specific saccharides. Besides it is involved in the development of acute and

chronic inflammation of the lung. Several human lung diseases are

characterized by decreased levels of bronchoalveolar SP-D.

Protein Information

Name SFTPD

Synonyms COLEC7, PSPD, SFTP4

Function Contributes to the lung's defense against inhaled microorganisms, organic

antigens and toxins. Interacts with compounds such as bacterial lipopolysaccharides, oligosaccharides and fatty acids and modulates leukocyte action in immune response. May participate in the extracellular reorganization or turnover of pulmonary surfactant. Binds strongly maltose

residues and to a lesser extent other alpha-glucosyl moieties.

Cellular Location Secreted, extracellular space, extracellular matrix. Secreted, extracellular

space, surface film

Tissue Location Expressed in lung, brain, pancreas and adipose tissue (mainly mature

adipocytes).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.