

# CD19

Catalog # PVGS1934

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">P25918</a> Mouse
<b>Sequence</b>	Arg19-Gly287
<b>Purity</b>	> 90% as determined by Bis-Tris PAGE
<b>Endotoxin Level</b>	Less than 1EU per $\mu$ g by the LAL method.
<b>Expression System</b>	HEK293
<b>Theoretical Molecular Weight</b>	30.7 kDa
<b>Formulation Reconstitution</b>	Lyophilized from a 0.22 $\mu$ m filtered solution in PBS, (pH 7.4). Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage &amp; Stability</b>	Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Gene ID</b>	12478
<b>Other Names</b>	B-lymphocyte antigen CD19, Differentiation antigen CD19, CD19, Cd19
<b>Target Background</b>	CD19 is a 95 kDa transmembrane glycoprotein that plays a central role in B cell activation and humoral immune responses. Functions as coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes. Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens. Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca <sup>2</sup> stores.

## Protein Information

---

<b>Name</b>	Cd19
<b>Function</b>	Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes. Decreases the threshold for activation of downstream

signaling pathways and for triggering B-cell responses to antigens (By similarity). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed:[12387743](#), PubMed:[20101619](#), PubMed:[9382888](#)). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:[7542548](#), PubMed:[7543183](#), PubMed:[9317126](#)). Required for normal differentiation of B-1 cells (PubMed:[12387743](#), PubMed:[7542548](#), PubMed:[7543183](#)). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:[12387743](#), PubMed:[7542548](#), PubMed:[9317126](#)). Required for normal levels of serum immunoglobulins, and for production of high- affinity antibodies in response to antigen challenge (PubMed:[12387743](#), PubMed:[7542548](#), PubMed:[7543183](#)).

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein

**Tissue Location**

Detected on B cells in spleen, bone marrow, thymus and lymph nodes (PubMed:[12387743](#), PubMed:[20101619](#), PubMed:[7542548](#)) Detected on peripheral blood lymphocytes (at protein level) (PubMed:[7543183](#)).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.